

CAPS Datasheets provide pest-specific information to support planning and completing early detection surveys.

Amrasca biguttula

Scientific Name

Amrasca biguttula (Ishida, 1912)

Synonym(s):

Amrasca biguttula biguttula (Ishida);
Amrasca devastans (Distant, 1918);
Chlorita biguttula Ishida; *Empoasca*
devastans Distant, 1918

Common Name

two-spot cotton leafhopper, cotton jassid, Indian green jassid, okra leafhopper

Type of Pest

Leafhopper

Taxonomic Position

Class: Insecta, **Order:** Hemiptera,
Family: Cicadellidae

Pest Recognition

This section describes characteristics of the organism and symptoms that will help surveyors recognize possible infestations in the field, select survey sites, and collect symptomatic material. For descriptions of diagnostic features, see the Identification/Diagnostic resources on the AMPS pest page on the CAPS Resource and Collaboration website.

Pest Description

Adults

Adult two-spot cotton leafhoppers are roughly 2-3 mm long and pale green with yellowish-green wings. Males are slightly smaller than females. This insect can be distinguished from most native cicadellid leafhopper species in North America by the pair of black spots on the head and the black spot at the apical end of each wing (Figs. 1 & 2). These markings may sometimes be faint or absent. When disturbed, adults and nymphs display characteristic sideways movements and hopping behaviors (Schreiner,



Figure 1: *Amrasca biguttula* adult (adapted from Cabrera-Asencio et al., 2023, [CC BY-NC 3.0](#))

2000). Confirmatory morphological identification requires dissection of adult male genitalia.



Figure 2: Dorsal view (A) and lateral view (B) of live *A. biguttula* adults on okra leaves from Florida. Note the blue arrows denoting the black spots on the head. Photos courtesy of Daphne Zapsas.

Eggs

Female two-spot cotton leafhoppers lay their eggs within the veins and midrib of host plant leaves, which renders the eggs nearly invisible without specialized tools (Mensah, 2006; Schreiner, 2000). The eggs are yellowish-white, oblong, and less than 0.04 inches long.

Nymphs

Newly hatched nymphs are wingless, pale green, and highly agile, quickly scuttling sideways or falling off of plants when disturbed (Schreiner, 2000). They are very small, usually less than 0.05 inches long (Fig. 3) (Jayasimha et al., 2012; Madar and Katti, 2011; Mensah, 2006) and are difficult to distinguish from the nymphs of other species. Like adults, they feed on the underside of leaves, stems, and petioles, causing damage and excreting sticky honeydew that further compromises plant health (AVRDC, 2003; Pestnet, 2021; Rajendran et al., 2018).

Nymphs coexist with adults throughout the year (AVRDC, 2003; Ghosh and Karmakar, 2022).




Figure 3: Nymphs and adults on okra leaves in Florida. Note nymphs clustering along the midrib of the leaves (**A**), an adult with prominent black spots on the wings (**B**), and possible "hopperburn", or discolored/necrotic tissue (**C**). Photo courtesy of Amy Roda.

Signs

Two-spot cotton leafhoppers cause significant damage to host plants resulting in disease-like symptoms collectively referred to as "hopperburn" (Fig. 4) (Rajendran et al., 2018; Schreiner, 2000; Schreiner, 1990). Both adults and nymphs feed on the underside of leaves, using their sucking mouthparts to consume the leaf cell contents and disrupt phloem tubes, which exacerbates the damage (Schreiner, 2000). Symptoms begin with yellow blotches on leaves, which progress to curling, browning, and dead spots, often accompanied by a distinctive yellow "halo" along the leaf edges (AVRDC, 2003; Schreiner, 2000). Severe infestations lead to bronzing, brittleness, defoliation, and stunted growth, with premature dropping of leaves, buds, and flowers significantly reducing plant productivity (Dung et al., 2021; Ghosh and Karmakar, 2022).



Figure 4: Damage to okra leaves by *A. biguttula* showing yellowing and some hopperburn (**A**), moderate hopperburn with leaf curling (**B**), and severe hopperburn with completely dead/dry portions of the leaf (**C**). Photos courtesy of Amy Roda.

Feeding also produces honeydew, which fosters the growth of sooty mold—a fungal layer that impedes photosynthesis and diminishes plant vitality (Kamble and Sathe, 2015; Rajendran et al., 2018).

Yield losses are particularly significant in cotton, okra, and sunflower (Ahmad et al., 1985; Al-Hamdany and Al-Karboli, 2017; Madar and Katti, 2011; Manivannan et al., 2021), though crops like eggplant and grapes can also be affected to a lesser extent (Ghosh and Karmakar, 2021; Lit and Bernardo, 1990; Schreiner, 1990).

Easily Mistaken Species

When surveying for *Amrasca biguttula* in the United States, expect to encounter many species of similar looking leafhoppers. They are often small and green, but most species lack the combination of a black spot on the forewing, two black spots on the head, as well as additional pattern elements. However, dissection of the male genitalia or DNA sequencing is necessary for positive morphological identification.

Here are a few domestic species that are similar in appearance and could be mistaken for *A. biguttula*:

Empoasca spp.

This genus includes multiple small, green leafhoppers that are about 1/8 inches in length. Nymphs are nearly identical to *A. biguttula* (Fig. 5 A). Adults are similar in size and color to *A. biguttula*; however, *Empoasca* spp. lack the distinct black spots on the anterior margin of the head and forewings (Fig. 5 B). Common species include *E. fabae* (Potato Leafhopper), *E. vitis* (Grape Leafhopper), and *E. kraemeri* (Bean Leafhopper).



Figure 5. *Empoasca fabae* nymph (A), and adult (B). Photo credits: (A) Frank Peairs, Colorado State University, [CC BY-NC 3.0](https://creativecommons.org/licenses/by-nc/3.0/) (B) Robert Webster / xpda.com, [CC-BY-SA-4.0](https://creativecommons.org/licenses/by-sa/4.0/)

Typhlocyba pomaria (White Apple Leafhopper)

This species is light-green and a similar size; however, it is more pale, yellowish white, and lacks the greenish tint and black spots (Fig. 6).




Figure 6. *Typhlocyba pomaria* (A) nymph and (B) adult. Photo credits: (A) University of Georgia Plant Pathology, University of Georgia, Bugwood.org [CC BY-NC 3.0](https://creativecommons.org/licenses/by-nc/3.0/), (B) shared from [iNaturalist](https://www.inaturalist.org/) © solomon v. hendrix, [CC BY-NC 4.0](https://creativecommons.org/licenses/by-nc/4.0/)

Kyboasca spp.

This is another genus of comparably sized, small, pale green leafhoppers with a similar black spot on each forewing (Fig. 7) but lacking black spots on the anterior margin of the head. It occurs in temperate areas of North America (Cabrera-Asencio et al., 2023; Dmitriev, (2022 onward)). This genus could be transient on *A. biguttula* host plants, but is primarily found on trees (Kittelberger and Howard, 2024).




Figure 7. Adults of *Kyboasca* spp. shared from [iNaturalist](#). Note the similar black spots on each forewing. Photo credits: (A) © Jason M Crockwell [CC BY-NC-ND 4.0](#) (B) © Melinda Fawver [CC BY-NC 4.0](#)

Alconeura spp.

This genus of leafhoppers has a similar spot on the forewings (Fig. 8), but the location is slightly different. It does not have black spots on the anterior margin of the head, and the insects often have red or orange coloration in addition to a green body (Cabrera-Asencio et al., 2023). The hosts of this genus are not well described, but there are no reports of *Alconeura* spp. on the hosts of *Amrasca biguttula*. This genus could be transient on *A. biguttula* host plants.



Figure 8. *Alconeura* spp. shared from [iNaturalist](#). Note the similar black spot on each forewing and the orange/yellow coloration. Photo credit: © Rob Van Epps [CC BY-NC 4.0](#).

Due to the similarities between *Amrasca biguttula* and other leafhoppers, morphological examination of male genitalia or molecular characterization is needed to confirm identification (Reddy et al., 2020; Sagarbarria et al., 2020).

Commonly Encountered Non-targets

Because the approved survey methods (visual surveys and trapping with yellow or green sticky traps) are non-specific, they will capture many non-target species, including the easily mistaken species described in the previous section.

There are currently no known lures or traps specific to this species.

Biology and Ecology

Amrasca biguttula (two-spot cotton leafhopper) females lay eggs on the underside of leaves or stems that hatch after approximately one week. Females are reported to lay between 14 and 60 eggs in their lifetime and eggs are typically laid within the veins and midribs of host plant leaves, rendering them nearly invisible without specialized tools (Jayasimha et al., 2012; Mensah, 2006; Schreiner, 2000).

Nymphs and adults live and feed together on the undersides of leaves (Dung et al., 2021; Schreiner, 2000), causing damage by sucking sap from mesophyll cells, resulting in curling, yellowing, and bronzing of leaves, a condition referred to as "hopperburn" (Kamble and Sathe, 2015; Reddy et al., 2020). Two-spot cotton leafhopper behavior on host plants is similar to other leafhoppers, including clustering on the underside of leaves, crawling, jumping, or flying short distances when disturbed, and secreting honeydew (Kamble and Sathe, 2015; Schreiner, 2000).

Reproduction occurs year-round, with overlapping generations, and adults living for about one month (Ghosh and Karmakar, 2022; Mensah, 2006). The lifecycle duration of the two-spot cotton leafhopper varies depending on climate and humidity. In India, generation times range from 15 to 46 days, with up to eleven generations per year (Ghosh and Karmakar, 2022; Mensah, 2006). With a continuous supply of host plants in warmer climates, two-spot cotton leafhoppers can breed and remain active year-round (AVRDC, 2003; Saeed et al., 2015). Population densities also fluctuate based on environmental conditions, increasing when temperatures are between 81°F and 93°F and relative humidity is around 70% and decreasing during periods of high rainfall and humidity (Ghosh and Karmakar, 2021; Mohammad et al., 2019). Populations typically peak during the hottest months of the year (Ahmad et al., 1985; Ghosh and Karmakar, 2022; Jayasimha et al., 2012; Mensah, 2006; Saeed et al., 2015; Thirasack, 2001).

Plant species attacked by the two-spot cotton leafhopper may shift throughout the season as the insects migrate to take advantage of available hosts (Ghosh and Karmakar, 2021; Saeed et al., 2015).

Known Hosts

Although the two-spot cotton leafhopper is polyphagous (Saeed et al., 2015), and a wide range of hosts are reported, most research focuses on incidence, damage, and control on cotton, okra, eggplant, and sunflower (Table 1) (Al-Hamdany and Al-Karboli, 2017; Fatima et al., 2021; Kalyan et al., 2017; Kamble and Sathe, 2015; Sahito et al., 2017). Grape is mentioned as a heavily affected susceptible crop in Ghosh and Karmakar (2021), and Kulkarni (2020) notes it as an emerging pest of grapevine in Maharashtra, India. However, we currently consider grapes a minor host due to limited evidence of impacts.

While not preferred targets for survey, Table 2 includes additional confirmed hosts of this species that can be a target for survey based on the needs of the surveyor.

The host lists below include cultivated and wild plants that 1) are infested by the pest under natural conditions, 2) are frequently described as major, primary, or preferred hosts, and 3) have primary evidence for feeding and damage documented in the literature. Plants are highlighted in bold if they are commercially produced and the pest causes economically significant damage.

Table 1. Preferred US hosts of *Amblyopoda biguttula*.

Scientific Name	Common Name	Type/Use	References/Notes
<i>Helianthus annuus</i> Linn.	Sunflower*	Cultivated	Kamble and Sathe, 2015; Saeed et al., 2015
<i>Abelmoschus esculentus</i> L.	Okra	Cultivated	Saeed et al., 2015
<i>Gossypium arboreum</i>	Cotton	Cultivated	Kamble and Sathe, 2015
<i>Gossypium hirsutum</i> L.	Cotton	Cultivated	Kamble and Sathe, 2015
<i>Solanum melongena</i> L.	Eggplant/brinjal	Cultivated	Saeed et al., 2015

* The majority of U.S. sunflower commercial production occurs in states with unsuitable climate conditions for *Amblyopoda* establishment

Table 2. Additional hosts of *Amrasca biguttula*.

Scientific Name (Common Name)	References
<i>Abutilon indicum</i> (L.) Sweet (monkeybush)	Saeed et al., 2015
<i>Achyranthes aspera</i> L. (devil's horsewhip)	Saeed et al., 2015
<i>Cajanus cajan</i> (L.) Millsp. (pigeon pea)	Kamble and Sathe, 2015
<i>Carthamus tinctorius</i> L. (safflower)	Javed et al., 2013
<i>Chenopodium murale</i> L. (nettleleaf goosefoot)	Saeed et al., 2015
<i>Citrullus lanatus</i> (Thumb) Mansf. (watermelon)	Saeed et al., 2015
<i>Cordia dichotoma</i> G. Forst (fragrant manjack)	Saeed et al., 2015
<i>Cucumis melo</i> L. var. phut (phut)	Saeed et al., 2015
<i>Cucumis melo</i> L. sativus (muskmelon)	Saeed et al., 2015
<i>Cucumis sativus</i> L. (cucumber)	Saeed et al., 2015
<i>Cyamopsis tetragonoloba</i> L. (guar)	Saeed et al., 2015
<i>Cynodon dactylon</i> (L.) Pers. (Bermuda grass)	Kamble and Sathe, 2015
<i>Datura innoxia</i> Mill. (D. metel L.) (pricklyburr)	Saeed et al., 2015
<i>Durio zibethinus</i> L. (durian)	Dung et al., 2021
<i>Grewia asiatica</i> L. (phalsa)	Saeed et al., 2015
<i>Guizotia abyssinica</i> (L.f.) Cass. (niger)	Darandale et al., 2015
<i>Hibiscus cannabinus</i> L. (mesta)	Raju and Rao, 1985
<i>Hibiscus rosa-sinensis</i> L. (China rose)	Kamble and Sathe, 2015
<i>Hibiscus sabdariffa</i> L. (mesta)	Raju and Rao, 1985
<i>Lagenaria siceraria</i> (Molina) Standl. (L. vulgaris Ser.) (bottle gourd)	Saeed et al., 2015

Scientific Name (Common Name)	References
<i>Luffa aegyptiaca</i> Mill. (sponge gourd)	Saeed et al., 2015
<i>Malva parviflora</i> L. (mallow)	Al-Hamdany and Al-Karbolli, 2017
<i>Nicotiana tabacum</i> L. (common tobacco)	Saeed et al., 2015
<i>Pisum sativum</i> L. (peas)	Saeed et al., 2015
<i>Raphanus sativus</i> L. (radish)	Navasero and Calumpang, 2013
<i>Ricinus communis</i> L. (castorbean)	Ishikawa and Takahata, 2019
<i>Sesamum orientale</i> L. (S. indicum L.) (sesame)	Ishikawa and Takahata, 2019
<i>Solanum inacum</i> Dunal (nightshade)	Ishikawa and Takahata, 2019
<i>Solanum muricatum</i> Ait. (pepino)	Ishikawa and Takahata, 2019
<i>Solanum tuberosum</i> L. (potato)	Kumar et al., 2011; Saeed et al., 2015
<i>Vigna mungo</i> (L.) Hepper (<i>Phaseolus mungo</i> L.) (black gram)	Saeed et al., 2015
<i>Vigna radiata</i> (L.) R. Wilczek (mung bean)	Navasero and Calumpang, 2013
<i>Vigna unguiculata</i> (L.) Walp. (cowpea)	Kamble and Sathe, 2015
<i>Vigna unguiculata</i> x <i>V. unguiculata</i> var. <i>sesquipedalis</i> (bush sitao)	Kamble and Sathe, 2015
<i>Vitis vinifera</i> L. (grapevine)	Ghosh and Karmakar, 2021
<i>Xanthium strumarium</i> L. (cocklebur)	Saeed et al., 2015

Pest Importance

Two-spot cotton leafhopper is a major agricultural pest that uses its piercing-sucking mouthparts to extract sap from leaves, causing hopperburn, a condition marked by yellowing, curling, and browning of leaves often followed by defoliation. This damage stunts plant growth and results in the premature drop of buds, flowers, and fruitlets, severely reducing crop yield and quality (Dung et al., 2021; Rajendran et al., 2018). Infestations can lead to yield losses ranging from 40% to 75% in key crops like cotton, okra, eggplant, and sunflower (Ahmad et al., 1985; Bhushan, 2023; Thapa et al., 2019).

The pest is particularly problematic in early growth stages of crops, affecting plants before traditional pest control measures are implemented (Navasero, 2015). In some cases, hopperburn increases the host plant's suitability to other pests, resulting in a complex of pests such as borers, thrips, and whiteflies (Sagarbarria et al., 2020). Beyond direct yield losses, infestations increase production costs as growers must adopt earlier and more frequent control measures, often relying on chemical pesticides (Ghosh and Karmakar, 2021).

Although *A. biguttula* does not typically kill host plants or directly affect natural ecosystems, its polyphagous nature means it feeds on a wide variety of plants, which could include plants federally listed as threatened or endangered. However, there is no evidence of overarching direct impacts on natural environments to date.

Market losses due to *A. biguttula* can be severe. In South Asia, it is a leading cause of reduced production in cotton, okra, and eggplant (Ghosh and Karmakar, 2021; Saeed et al., 2015). In Africa, recent infestations have contributed to a 38% reduction in cotton exports from Senegal and significant financial losses for farmers, prompting government subsidies and loan forgiveness programs (PR-PICA, 2022; Sylla, 2023). Detection of *A. biguttula* is challenging due to inefficient survey methods and resemblance to other leafhopper species, increasing the likelihood of unnoticed establishment and subsequent economic impacts (Navasero, 2015)

Potential trade implications could arise if infestations establish in new regions like the United States, where crops such as cotton, eggplant, and grapes are economically important.

Pathogens or Associated Organisms Vected

Amrasca biguttula is not confirmed to be a vector of any pathogens.

This species has been associated with phytoplasma species and chickpea chlorotic dwarf virus, but its status as a vector has not been confirmed (Biswas et al., 2018; Nielson, 1968; Reddy et al., 2020; Sharif et al., 2019).

Known Distribution

The two-spot cotton leafhopper is native to Asia and Oceania, where it is widespread in warmer areas, occurring from the Middle East to Australia and the Mariana Islands. However, recent outbreaks of the two-spot cotton leafhopper in Africa (PR-PICA, 2022) and new detections in the Caribbean (Cabrera-Asencio et al., 2023; IPPC, 2024) suggest this pest is spreading rapidly. The two-spot cotton leafhopper may have been present in Africa at non-outbreak levels for some time, with detections in Ghana as early as 1999 (Obeng-Ofori and Sackey, 2003), but the pest appears to have spread recently into new regions across Africa and it was reported for the first time in Northern Cameroon in 2022 (Jacques et al., 2024).

Table 3. Countries where *A. biguttula* is known to occur.

Continent	Country	References/Notes
Africa	Benin	PR-PICA, 2022
Africa	Burkina Faso	PR-PICA, 2022
Africa	Cameroon	Jacques et al., 2024; PR-PICA, 2022
Africa	Chad	PR-PICA, 2022
Africa	Côte d'Ivoire	PR-PICA, 2022
Africa	Ghana	Obeng-Ofori and Sackey, 2003
Africa	Mali	PR-PICA, 2022
Africa	Niger	Franç et al., 2024
Africa	Senegal	PR-PICA, 2022
Africa	Togo	PR-PICA, 2022

Asia	Afghanistan	Xu et al., 2017
Asia	Bangladesh	Xu et al., 2017
Asia	China	Xu et al., 2017
Asia	India	Xu et al., 2017
Asia	Indonesia	Xu et al., 2017
Asia	Iraq	Al-Hamdany and Al-Karboli, 2017
Asia	Japan	Thirasack, 2001
Asia	Laos	Thirasack, 2001
Asia	Pakistan	Xu et al., 2017
Asia	Philippines	Xu et al., 2017
Asia	Sri Lanka	Xu et al., 2017
Asia	Thailand	Xu et al., 2017
Asia	Vietnam	Xu et al., 2017
North America	Barbados	IPPC, 2024
Oceania	Australia (Christmas Island)	Kulkarni, 2020
Oceania	Guam	Schreiner, 1990
Oceania	Northern Mariana Islands	Schreiner, 1990

Status of infestation in the United States (January 2025)

In the Caribbean, the pest was identified in the southern portion of Puerto Rico in 2023 (Cabrera-Asencio et al., 2023) and the U.S. Virgin Islands in 2024 (FAO, 2022; IPPC, 2024). In November 2024, two-spot cotton leafhopper was detected in Florida.

Pathway

The movement of *Amrasca biguttula* into new regions has not been linked to any specific pathway. Recent spread and new introductions of *A. biguttula* indicate there is an unknown open pathway for this pest to move to new areas. Agricultural inspectors have intercepted *A. biguttula* five times from aircraft cargo at ports of entry in the United States (ARM, 2024), suggesting that, while it appears to happen rarely, the two-spot cotton leafhopper is capable of moving into the country through international trade. Adults and nymphs have the potential to travel as hitchhikers on humans, equipment, or other transport mechanisms and eggs laid on plants imported for propagation could introduce the pest to new regions.

Natural spread may also occur, though adults are unlikely to disperse long distances unaided based on the short dispersal distance of similar leafhopper species (Blackmer et al., 2004; Northfield et al., 2009). However, there have been documented cases of leafhoppers being carried on wind currents over extremely long distances (Ghauri, 1982) and there is potential for wind-assisted long distance dispersal from the Caribbean into the continental United States (Andraca-Gómez et al., 2020; Bertone et al., 2008).

Use the PPQ Commodity Import and Export manuals listed below to determine 1) if host plants or material are allowed to enter the United States from countries where the

organism is present and 2) what phytosanitary measures (e.g., inspections, phytosanitary certificates, post entry quarantines, mandatory treatments) are in use. These manuals are updated regularly.

[Agricultural Commodity Import Requirements \(ACIR\) search tool](#): A single source to search for and retrieve entry requirements for imported commodities.

[Plants for Planting Manual](#): This manual is a resource for regulating imported plants or plant parts for propagation, including buds, bulbs, corms, cuttings, layers, pollen, scions, seeds, tissue, tubers, and like structures.

[Treatment Manual](#): This manual provides information about treatments applied to imported and domestic commodities to limit the movement of agricultural pests into or within the United States.

Potential Distribution within the United States

Based on a comparison of plant hardiness zones (PHZ) (Takeuchi et al., 2018) where this species is known to occur and collection records exist, it could establish in the United States in PHZ 7-14. This area spans the eastern seaboard from Massachusetts to Florida, west through most of Virginia, southern parts of Kentucky, Missouri, Kansas, Colorado, Utah, and Nevada. It also extends across the southern states, along the west coast from the Mexican border to the Canadian border, and to Hawaii and the territories.

Areas in the United States in PHZ 10-12 may be at greater risk for establishment and impacts based on more frequent distribution records in those climates. The distribution of hosts is unlikely to significantly affect the area at risk, as the species is a polyphagous pest that regularly changes host plants based on availability.

Survey and Key Diagnostics

Approved Methods for Pest Surveillance*:

For the current approved methods and guidance for survey and identification, see [Approved Methods for Pest Surveillance \(AMPS\)](#) pest page on the CAPS Resource and Collaboration website.

Survey Method images

See below for images of the recommended visual survey/manual collection methods. The full description of survey design guidance and sample collection instructions are found at the [approved methods site](#).



Figure 9. Bagging host leaves to collect specimens. **(A)** Inverting the bag over the leaf and **(B)** collecting the entire leaf to capture all insects found. Photos courtesy of Daphne Zapsas (A) and Amy Roda (B).



Figure 10. Beat sampling by tapping host plant leaves over a tray filled with alcohol. Photo courtesy of Daphne Zapsas.

References

- Ahmad, Z., Attique, M.R. and Rashid, A., 1985. An estimate of the loss in cotton yield in Pakistan attributable to the jassid *Amrasca devastans* Dist. Crop Protection, 5(2): 105-108.
- Al-Hamdany, M.N. and Al-Karboli, H.H., 2017. First record of okra leafhopper, *Amrasca biguttula biguttula* Ishida on okra in Iraq. International Journal of Agricultural Technology, 13(3): 393-402.
- Andraca-Gómez, G., Lombaert, E., Ordano, M., Pérez-Ishiwara, R., Boege, K., Domínguez, C.A. and Fornoni, J., 2020. Local dispersal pathways during the invasion of the cactus moth, *Cactoblastis cactorum*, within North America and the Caribbean. Scientific Reports, 10(1): 11012.
- ARM, 2024. Agricultural Risk Management (ARM) System. United States Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine.
- Atakan, E., 2011. Development of a sampling strategy for the leafhopper complex [*Asymmetrasca decedens* (Paoli) and *Empoasca decipiens* Paoli] (Hemiptera: Cicadellidae) in cotton. Journal of Pest Science, 84(2): 143-152.
- AVRDC, 2003. A farmer's guide to harmful and helpful insects in eggplant fields. Asian Vegetable Research and Development Center (AVRDC), The World Vegetable Center, Shanhua, Taiwan, 22 pp.
- Bertone, C.A., Meissner, H.E. and Lemay, A.V., 2008. Natural spread of pests within and into the greater Caribbean region, Caribbean Food Crops Society, 44th Annual Meeting, Miami, Florida, USA.
- Bhushan, L.S., Raut, A.M., Talekar, N., Chaudhary, B., Sonawane, V., and Ashwini, A., 2023. Biopesticides and their impact on brinjal Jassid (*Amrasca biguttula biguttula* Ishida). Journal of Applied and Natural Science, 15(4): 1398-1406.
- Biswas, C., Dey, P., Meena, P.N., Satpathy, S. and Sarkar, S.K., 2018. First report of a subgroup 16SrIX-E ('*Candidatus* Phytoplasma phoenicium'-related) phytoplasma associated with phyllody and stem fasciation of sunn hemp (*Crotalaria juncea* L.) in India. Plant Disease, 102(7): 1445.
- Blackmer, J.L., Hagler, J.R., Simmons, G.S. and Cañas, L.A., 2004. Comparative dispersal of *Homalodisca coagulata* and *Homalodisca liturata* (Homoptera: Cicadellidae). Environmental Entomology, 33(1): 88-99.
- Cabrera-Asencio, I., Dietrich, C.H. and Zahniser, J.N., 2023. A new invasive pest in the Western Hemisphere: *Amrasca biguttula* (Hemiptera: Cicadellidae). Florida Entomologist, 106(4): 263-266.
- Dmitriev, D.A., Angelova, R., Anufriev, G.A., Bartlett, C.R., Blanco-Rodríguez, E., Borodin, O.I., Cao, Y.-H., Deitz, L.L., Dietrich, C.H., Dmitrieva, M.O., El-Sonbati, S.A., Evangelista de Souza, O., Gjonov, I.V., Gonçalves, A.C., Gonçalves, C.C., Hendrix, S., McKamey, S., Kohler, M., Kunz, G., Malenovský, I., Morris, B.O., Novoselova, M., Pinedo-Escatell, J.A., Rakitov, R.A., Rothschild, M.J., Sanborn, A.F., Takiya, D.M., Wallace, M.S., Zahniser, J.N., (2022 onward). *Kyboasca* Zakhvatkin, 1953. World Auchenorrhyncha Database. TaxonPages.
- Dung, L.T., Liem, N.V., Lam, P.V., Tan, L.V. and Hai, N.N., 2021. Morphological and molecular characterization of two leafhopper species of genus *Amrasca* (Hemiptera: Cicadellidae) hosting to durian. Journal of Entomological Research,

- 45(3): 476-481.
- FAO, 2022. GIAHS Proposal: Qingyuan Forest-Mushroom Co-culture System in Zhejiang Province, Food and Agriculture Organization of the United Nations.
- Fatima, T., Srinivas, P.S., Sridevi, G., Shivani, D. and Bharati, B., 2021. Biochemical responses in resistant and susceptible sunflower accessions elicited by leafhopper, *Amrasca biguttula biguttula* Ishida feeding. *Journal of Experimental Zoology India*, 24: 1331-1336.
- Franç, Z., Moussa, O.Z., Atta, S., Leyo, I.H. and Guimbo, I.D., 2024. Cotton leafhoppers, *Amrasca biguttula* (Ishida, 1913)(Hemiptera: Cicadellidae), identified as a new species on okra and guinea sorrel in Niger. *Advances in Entomology*, 12(3): 183-194.
- Ghauri, M.S.K., 1982. A case of long distance dispersal of a leafhopper. In: T.S. Robertson, M.R. Wilson, W.J. Knight and N.C.Pant (Editors), 1st International Workshop on Leafhoppers and Planthoppers of Economic Importance. Commonwealth Institute of Entomology, London, UK, pp. 249-255.
- Ghosh, S.K. and Karmakar, R., 2021. Sustainable control of leaf hopper (*Amrasca biguttula biguttula* Ishida) on grape vine (*Vitis vinifera* L.). *Journal of Zoology*, 42(21): 83-99.
- Ghosh, S.K. and Karmakar, R., 2022. Sustainable management of leaf hopper (*Amrasca biguttula biguttula*) on eggplant/brinjal (*Solanum melongena* (Linn.) and related crops. In: W. Hasan (Editor), *Hi-tech Crop Production and Pest Management*. Biotech Books, pp. 233-258.
- IPPC, 2024. First report of presence of Indian cotton leafhopper (*Amrasca biguttula*), The International Plant Protection Convention (IPPC), Rome, Italy.
- Jacques, H.D., Socrates, D.N., Bouladji, Y., Djague, T.L., Moïse, A. and Noé, W., 2024. First infestation of an exotic crops pest, *Amrasca biguttula* on cotton *Gossypium hirsutum* L. in North Cameroon. *International Journal of Plant & Soil Science*, 36(3): 16-22.
- Jayasimha, G.T., Rachana, R.R., Manjunatha, M. and Rajkumar, V.B., 2012. Biology and seasonal incidence of leafhopper, *Amrasca biguttula biguttula* (Ishida) (Hemiptera: Cicadellidae) on okra. *Pest Management in Horticultural Ecosystems*, 18(2): 149-153.
- Kalyan, R., Saini, D., Meena, B., Pareek, A., Naruka, P., Verma, S. and Joshi, S., 2017. Evaluation of new molecules against jassids and whiteflies of Bt cotton. *Journal of Entomology and Zoology Studies*, 5(3): 236-240.
- Kamble, C.S. and Sathe, T.V., 2015. Incidence and host plants for *Amrasca biguttula* (Ishida) from Kolhapur region, India. *International Journal of Development Research*, 5(3): 3658-3661.
- Khaing, O., Hormchan, P., Jamornmarn, S., Ratanadilok, N. and Wongpiyasatid, A., 2002. Spatial distribution pattern of cotton leafhopper, *Amrasca biguttula* (Ishida)(Homoptera: Cicadellidae). *Kasetsart Journal (Natural Science)*, 36: 11-17.
- Kittelberger, K. and Howard, T., 2024. Hoppers of North Carolina: Spittlebugs, Leafhoppers, Treehoppers, and Planthoppers. North Carolina Biodiversity Project and North Carolina State Parks, Raleigh, NC.
- Kulkarni, N., 2020. Sucking pests of grapes. In: Omkar (Editor), *Sucking Pests of Crops*.

- Springer Singapore, Singapore, pp. 425-450.
- Lashari, A.A., Korai, S.K., Nizamani, I.A., Qureshi, K.H., Lodhi, A.M., Korai, A.K. and Korai, P.K., 2022. Monitoring of sucking pest on mustard crop through different colours sticky traps. *Pakistan Journal of Zoology*, 54(2): 810-808.
- Lit, M.C. and Bernardo, E.N., 1990. Mechanism of resistance of eggplant (*Solanum melongena* Linn.) to the cotton leathopper, *Amrasca biguttula* (Ishida) II. morphological and biochemical factors associated with resistance. *Philippine Journal of Crop Science*, 15(2): 79-84.
- Madar, H. and Katti, P., 2011. Biology of leafhopper, *Amrasca biguttula biguttula* on sunflower. *International Journal of Plant Protection* 2(4): 370-373.
- Madhu, T.N., Kumar, R., Naik, V.C., Prabhulinga, T., Santosh, S. and Chandrashekar, N., 2022. Attraction of leaf hopper, *Amrasca biguttula biguttula*, and whitefly, *Bemisia tabaci*, toward natural essential oils in cotton. *Animal Biology*, 73(1): 27-41.
- Manivannan, A., Kanjana, D., Dharajothi, B. and Meena, B., 2021. Evaluation of resistance in cotton genotypes against leafhoppers *Amrasca biguttula biguttula* (Ishida), (Homoptera: Cicadellidae). *International Journal of Tropical Insect Science*, 41: 2409-2420.
- Mensah, R.K., 2006. National Cotton Industry Biosecurity Plan: Appendix 3. Pest Risk Reviews, Plant Health Australia.
- Mohammad, A., Alam, S.N., Miah, M.R.U., Amin, M.R. and Smriti, R.S., 2019. Population fluctuation of jassid, and shoot and fruit borer of okra. *Bangladesh Journal of Agricultural Research*, 44(3): 493-499.
- Navasero, M., 2015. Insect pests of eggplant. *Pests and Diseases of Economically Important Crops in the Philippines*. Pest Management Council of the Philippine, Inc. c/o Crop Protection Cluster, University of the Philippines Los Banos, College, Laguna: 354-383.
- Nielson, M.W., 1968. The leafhopper vectors of phytopathogenic viruses (Homoptera, Cicadellidae): Taxonomy, biology and virus transmission., United States Department of Agriculture, Agricultural Research Service, Washington, D. C.
- Northfield, T.D., Mizell, R.F., III, Paini, D.R., Andersen, P.C., Brodbeck, B.V., Riddle, T.C. and Hunter, W.B., 2009. Dispersal, patch leaving, and distribution of *Homalodisca vitripennis* (Hemiptera: Cicadellidae). *Environmental Entomology*, 38(1): 183-191.
- Obeng-Ofori, D. and Sackey, J., 2003. Field evaluation of non-synthetic insecticides for the management of insect pests of okra *Abelmoschus esculentus* (L.) Moench in Ghana. *Ethiopian Journal of Science*, 26(2): 145-150.
- Pestnet, 2021. Pacific Pests, Pathogens & Weeds: Bele (*Abelmoschus*) leafhoppers (039). Australian Centre for International Agricultural Research.
- PR-PICA, 2022. PR-PICA newsletter No. 44, Oct - Nov 2022, Programme regional de production integree du coton en afrique (PR-PICA) [Regional integrated cotton production program in Africa], Bobo-Dioulasso, Burkina Faso.
- Rajendran, T.P., Birah, A. and Burange, P.S., 2018. Insect pests of cotton. In: Omkar (Editor), *Pests and Their Management*. Springer Nature, Singapore, pp. 361-411.
- Reddy, M.G., Rao, G.P. and Meshram, N.M., 2020. Molecular identification of leafhopper potential vectors of chickpea stunt using the COI sequences.

- Phytopathogenic Mollicutes, 10(2): 214-219.
- Saeed, R., Razaq, M. and Hardy, I.C.W., 2015. The importance of alternative host plants as reservoirs of the cotton leaf hopper, *Amrasca devastans*, and its natural enemies. *Journal of Pest Science*, 88: 517-531.
- Sagarbarria, M.G.S., Taylo, L.D. and Hautea, D.M., 2020. Morphological and molecular analyses of leafhopper, *Amrasca biguttula* (Ishida) (Hemiptera: Cicadellidae) infesting eggplant (*Solanum melongena* L.) in Luzon Island, Philippines. *Journal of Asian-Pacific Entomology*, 23: 260-268.
- Sahito, H.A., Shah, Z.H., Kousar, T., Mangrio, W.M., Mallah, N.A., Jatoi, F.A. and Kubar, W.A., 2017. Research article comparative efficacy of novel pesticides against Jassid, *Amrasca biguttula biguttula* (Ishida) on cotton crop under field conditions at Khairpur, Sindh, Pakistan. *Singapore Journal of Scientific Research*, 1: 1-8.
- Schreiner, I., 2000. Okra leafhopper (*Amrasca biguttula* Ishida), *Agricultural Development in the American Pacific*.
- Schreiner, I.H., 1990. Resistance and yield response to *Amrasca biguttula* (Homoptera: Cicadellidae) in eggplant (*Solanum melongena*). *Philippine Entomologist*, 8(1): 661-669.
- Sharif, M.Z., Ahmad, S.J.N., Tahir, M., Ziaf, K., Zhang, S.-H. and Ahmad, J.N., 2019. Molecular identification and characterization of phytoplasmas associated with carrot, cabbage and onion crops and their insect vectors in Punjab, Pakistan. *Pakistan Journal of Agricultural Sciences*, 56(2): 407-414.
- Sivakumar, G., Mohan. M, Subaharan, K., Venkatesan, T., Yelshetti, S., Kannan, M., Anandham, R., Yandigeri, M.S., Kumari, S., Elango, K. and Ram Kumar, P., 2022. Gut bacteria mediated insecticide resistance in cotton leafhopper *Amrasca biguttula biguttula*. *Current Science*, 122(8): 958-964.
- Srinivasan, K., Krishnakumar, N.K., Ramachander, P.R. and Rao, G.S.P., 1988. Seasonal patterns of leafhopper, *Amrasca biguttula biguttula* on okra in India. *International Journal of Tropical Insect Science*, 9(1): 85-88.
- Sylla, F., 2023. Cotton and products update, United States Department of Agriculture - Foreign Agricultural Service, Washington, D.C.
- Takeuchi, Y., Flower, G. and Joseph, A.S., 2018. SAFARIS: Global Plant Hardiness Zone Development. North Carolina State University, Center for Integrated Pest Management; United States Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine, Science and Technology, Plant Epidemiology and Risk Analysis Laboratory, Raleigh, N.C., pp. 6.
- Thapa, R., Bista, K., Bhatta, M., Bhandari, S., Acharya, S.R. and Sapkota, B., 2019. Comparative performance and economic efficiency of different pesticides against okra jassids (*Amrasca biguttula biguttula*): Their impact on okra yield and growth attributes. *Journal of Entomology and Zoology Studies*, 7(5): 525-531.
- Thirasack, S., 2001. Yield losses assessment due to pests on cotton in Lao PDR. *Agriculture and Natural Resources*, 35(3): 271-283.
- Wagen, T.A., Jessar, M.A., Abbasi, N.H., Khuhro, S., Abro, M.A., Bughio, A.N., Junejo, A.S. and Pathan, G.M., 2019. Evaluation of colored sticky traps for monitoring and managing the jassid population in okra. *Journal of Biology, Agriculture and Healthcare*, 9(11): 35-38.

Wille, H.P., 1950. Untersuchungen über *Psylla piri* L. und andere Birnblattsaugerarten im Wallis, ETH Zurich.

Xu, Y., Wang, Y., Dietrich, C.H., Fletcher, M.J. and Qin, D., 2017. Review of Chinese species of the leafhopper genus *Amrasca* Ghauri (Hemiptera, Cicadellidae, Typhlocybinae), with description of a new species, species checklist and notes on the identity of the Indian cotton leafhopper. *Zootaxa*, 4353(2): 360-370.

USDA-APHIS-PPQ-ST staff developed this datasheet. Cite this document as:

PPQ. 2025. Cooperative Agricultural Pest Survey (CAPS) Pest Datasheet for *Amrasca biguttula* (Cicadellidae): Two-spot cotton leafhopper. United States Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine (PPQ), Raleigh, NC.

Versions

January 2025: Datasheet completed (Version 1).

May 2026: Common name changed from cotton jassid to two-spot leafhopper (Version 2).

Reviewers

Nichole Carrier, National Operations Manager, USDA-APHIS-PPQ-FO

Todd Gilligan, Director, USDA-APHIS-PPQ-S&T, PITL

Stephen Lavalley, National Policy Manager, USDA-APHIS-PPQ-EDP, PER

Eric Rohrig, National Science Program Coordinator, USDA-APHIS-PPQ-S&T, DESS

Amy Roda, Supervisory Entomologist, USDA-APHIS-PPQ-S&T, TIML