CAPS Datasheets provide pest-specific information to support planning and completing early detection surveys.

Ralstonia solanacearum Race 3 Biovar 2

Scientific Name

Ralstonia solanacearum (Smith, 1896) Yabuuchi et al. (1995), phylotype IIB, sequevars 1 and 2 (known historically and for regulatory purposes as race 3 biovar 2)

Synonyms:

Burkholderia solanacearum (Smith,1896) Yabuuchi et al.,1993
Pseudomonas solanacearum (Smith,1896) Smith,1914

Common Names

Brown rot of potato, Southern wilt of geranium, bacterial wilt of tomato

Type of Pest

Bacterium

Taxonomic Position

Class: Betaproteobacteria, **Order:** Burkholderiales, **Family:** Burkholderiaceae

Notes on taxonomy and nomenclature:

Four phylotypes, corresponding to geographic origin and three species have been identified in the *R. solanacearum* species complex based on sequence analyses. *Ralstonia solanacearum* (phylotype II) originated in the Americas, *R. pseudosolanacearum* (phylotype I and III) originated in Asia and Africa, respectively, and *R. syzygii* (phylotype IV) originated in the Indonesian Archipelago (Prior et al., 2016; Safni et al., 2014). Strains in phylotype IIB sequevar 1-2 (abbreviated IIB-1 and IIB-2) are known for regulatory purposes as race 3 biovar 2 (EPPO, 2022b; USDA-APHIS-PPQ, 2022).

Pest Recognition

This section describes characteristics of the organism and symptoms that will help surveyors recognize possible infestations/infections in the field, select survey sites, and collect symptomatic material. For morphological descriptions, see the Identification/Diagnostic resources on the AMPS pest page on the CAPS Resource and Collaboration website.

Pest Description

This datasheet presents information for *Ralstonia solanacearum* Race 3 Biovar 2 (herein abbreviated R3 Bvr2), also known as phylotype IIB, sequevars 1 and 2. It is a quarantine pathogen in Europe and United States (EPPO, 2022b; Ozakman and Schaad, 2003).

Symptoms

Primary symptoms of bacterial wilt induced by R3 Bvr2 include leaf wilting (often as upward-rolling margins appearing unilaterally, leaf chlorosis (yellowing), necrosis (browning) of vascular tissue, stunting, vascular rings, and rotting of tubers (Champoiseau et al., 2009; Swanson et al., 2007; Swanson et al., 2005). Wilted leaves are usually the first symptom, followed by chlorosis and plant death. In the early stages of the disease, infected plants may appear to recover at night (Champoiseau et al., 2009). The stem may collapse, and gray-white bacterial ooze may be present, especially when the stem is cut or broken. Stems may also blacken (Janse, 2004). A common diagnostic sign of bacterial wilt is bacterial streaming (Fig. 1), which occurs when freshly cut stems from infected plants are placed in water. Fine, milky white strands of a viscous white slime containing bacteria often run from the cut end of the stem within 15 minutes (Dickstein et al., 2017).

On potatoes and other solanaceous hosts: The first visible symptom of *R. solanacearum* R3 Bvr2 on solanaceous hosts is wilting of the youngest leaves during the hottest part of the day, often on just one side of a leaflet or on a single branch (Fig. 2a). Another common field symptom associated with bacterial wilt is plant stunting. The entire plant may rapidly wilt under favorable conditions, starting with wilting and yellowing of foliage, and eventually leading to plant death. In infected potato tubers, the vascular ring turns a grey-brown color, which can spread into the pith or cortex as the infection advances. When these infected potatoes are sliced open, they release a milky-white sticky exudate. Threads may become apparent from the ooze when the two sides of a cut potato are pressed together and then separated. Additionally, ooze may emerge from the tuber eyes, causing dirt to stick on tuber surface (Champoiseau et al., 2009) (Fig. 2b). In tomato, the disease develops rapidly and plants may die within 4–7 days of symptoms appearing (Dickstein et al., 2017). Latently infected tomato plants may wilt suddenly after fruit set. A longitudinal slice of infected stems or stolons will reveal vascular browning, visible as long, narrow, dark brown streaks (Fig. 3). In succulent young plants of highly susceptible varieties, the stem can collapse, and gray-white bacterial ooze may be visible on stem surfaces (Fig. 3) (Champoiseau et al., 2009; Dickstein et al., 2017).



Figure 1: Shows positive results for bacterial streaming. Photo Courtesy of David B. Langston, University of Georgia, Bugwood.org.




Figure 2: Wilting potato plant and grey-brown vascular ring in a tuber caused by *R. solanacearum* R3 Bvr2. Image credits: Amilcar Sanchez Perez (a) and Caitilyn Allen, University of Wisconsin-Madison (b).



Figure 3: Generalized symptoms of tomato wilt caused by *R. solanacearum* R3 Bvr2 **Left:** Wilted tomato. **Right:** Vascular browning and brown streaks observed in a tomato stem. Image credits: Clemson University - USDA Cooperative Extension Slide Series (left) and Central Science Laboratory, Harpenden Archive, British Crown (right). All photos are from Bugwood.org.



Figure 4: Left: A geranium showing early wilting symptoms **Right:** A geranium showing yellowing symptoms caused by *R. solanacearum* R3 Bvr2. Image credits: Wisconsin Department of Agriculture, Trade and Consumer Protection (left) and Caitilyn Allen, University of Wisconsin-Madison (right).

In geranium: The common name for this disease in geranium is southern wilt. Symptoms can be subtle and easily overlooked. Under favorable conditions, the disease develops rapidly, and symptoms move upwards from older to younger leaves.

Chlorosis and wilting of lower leaves usually appear first, followed by upward curling of leaf margins, which is an identifying characteristic for the disease in geranium. The leaf margins may become chlorotic and then necrotic. Wilted leaves often develop wedge-shaped areas of chlorosis that also become necrotic (Fig. 4). Vascular discoloration is visible in stems (especially at the root crown) (Fig. 5) and in the roots; these areas can blacken and eventually become necrotic (Dickstein et al., 2017; Janse et al., 2004). In late stages of disease, stems may collapse and the whole plant may desiccate and die.

Easily Mistaken Species

Symptoms of *R. solanacearum* R3 Bvr2 infection may resemble those caused by other plant pathogens or abiotic stress. A diagnosis should not be based solely on symptoms (Dickstein et al., 2017).

Other *R. solanacearum* strains that are endemic in the southern United States produce similar symptoms in the same hosts (Dickstein et al., 2017). *Xanthomonas hortorum* pv. *pelargonii* causes similar wilting symptoms in geranium (Moorman, 2014). *Clavibacter michiganensis*



Figure 5: Bacterial ooze in geranium stem caused by infection with R. solanacearum R3Bv2. Image credits: Margery Daughtrey, Cornell University, Bugwood.org.

subsp. *sepedonicus* produces similar vascular browning in potato (Stevenson et al., 2001). *R. solanacearum* R3 Bvr2 symptoms on potato can also be confused with those caused by *Pectobacterium* spp., *Fusarium* spp., and *Verticillium* spp. (Cook and Sequeira, 1991). Abiotic stresses, such as drought, mechanical damage to the roots, or nutrient deficiency can also produce symptoms easily mistaken for *R. solanacearum* R3 Bvr2 (Dickstein et al., 2017).

Biology and Ecology

Ralstonia solanacearum R3 Bvr2 is a gram-negative soil- and waterborne vascular bacterial pathogen (Marco-Noales et al., 2008; Swanson et al., 2007; Wenneker et al., 1999). *R. solanacearum* R3 Bvr2 can live in the roots of hosts, including inside potato tubers, in the soil matrix surrounding the roots (rhizosphere), and in plant debris (Granada and Sequeira, 1983; Janse et al., 2004; Wenneker et al., 1999). The bacterium will grow on a variety of non-selective and semi-selective media between the temperatures of 70°F and 88°F (Caruso et al., 2003; Cook and Sequeira, 1991; Marco-Noales et al., 2008; Ozakman and Schaad, 2003; Mahbou Somo Toukam et al., 2009). *R. solanacearum* R3 Bvr2 grows slowly in culture and is easily outcompeted by other microbes; individual bacterial colonies that appear in less than 36 hours at 82°F are usually not *R. solanacearum* (Cook and Floyd, 2020).

Ralstonia solanacearum R3 Bvr2 can infect new hosts in many ways because it can survive in water, soil, and within infected plants for a long period of time. The pathogen can enter xylem tissue through wounds (Milling et al., 2009), spread via infected plant cuttings and seed tubers, and invade the plant through wounds created by nematodes or emerging roots (Swanson et al., 2007). When infected plants decompose, bacteria are released into the environment. Plants will also release the bacteria from open wounds. The bacteria emerging from the wound creates a gray-white bacterial ooze made from protective polysaccharides to help it survive in the environment (Shekhawat and Perombelon, 1991; van Elsas et al., 2000). Bacterial cells easily spread via irrigation and surface runoff water, infested soil, tools and equipment (Cook and Floyd, 2020; Swanson et al., 2007).

Unlike most strains of *Ralstonia solanacearum*, *R. solanacearum* R3 Bvr2 thrives in the cool tropical highlands and in temperate zones. This bacterium is highly virulent at temperatures between 66 ºF and 82 ºF (Huerta et al., 2015). Virulence decreases with temperatures above 95 ºF or below 61 ºF (Champoiseau et al., 2009; Ciampi and Sequeira, 1980).

R3 Bvr2 is also known to be more cold-tolerant than the southeastern native/endemic species (phylotype IIC, sequevar, commonly referred to as R1 Bvr1) and may survive throughout the continental United States (Milling et al., 2009; Swanson et al., 2007; Swanson et al., 2005). In Australia, R3 Bvr2 can survive in fallow soils with temperatures in winter as low as 39°F and in potato tubers at a constant cold storage temperature of 39°F for up to 4 months (Milling et al., 2009). However, laboratory studies found that R3 Bvr2 cannot survive the repeated freeze-thaw cycles that are common in northern continental U.S. states (Scherf et al., 2010).

Surveys for R3 Bvr2 are often difficult because the bacterium can evade detection in latently infected (symptomless) hosts (Swanson et al., 2005). Under temperate conditions, this pathogen survives in deep soil (~2 ft) (Graham and Lloyd, 1979; van Elsas et al., 2000) and is known to survive up to two years in soil where affected crops have already been removed (van Elsas et al., 2000; Wenneker et al., 1999). In warmer regions of South Africa, R3 Bvr2 can persist in the soil for more than 5 years even without known hosts present (Stander et al., 2003).

Known Hosts

The host list below includes cultivated and wild plants that 1) are infected or infested by the pest under natural conditions, 2) are frequently described as major, primary, or preferred hosts, and 3) have primary evidence for feeding and damage documented in the literature. Plants are highlighted in bold if they are commercially produced and the pest causes economically significant damage.

Economically important hosts of *R. solanacearum* R3 Bvr2 include potato, tomato, and geranium. It has also been known to colonize several weed species. These latent infections play a crucial role in the dissemination of *R. solanacearum* R3 Bvr2 and significantly influence the epidemiology of the disease (Dickstein et al., 2017). In Europe, numerous outbreaks of *R. solanacearum* R3 Bvr2 have been attributed to bittersweet nightshade (*Solanum dulcamara* L.), a weed species that grows along waterways and in wet areas in N. America and Europe (Janse, 1996; Parkinson et al., 2013). In addition to its primary economic hosts — potato, tomato, and geranium —R3 Bvr2 colonizes many other plant hosts. This adaptability poses a persistent threat to subsequent planting seasons and should be considered when designing a survey.

Major Hosts

Pelargonium hortorum* (zonal geranium), ***Pelargonium* spp. (geranium)* , *Pelargonium zonale* (horseshoe geranium)***, ***Solanum lycopersicum* (tomato)*,** and ***Solanum tuberosum* (potato)*** (Ciampi and Sequeira, 1980; Gutarra et al., 2017; Janse et al., 2004; Milling et al., 2009; Ozakman and Schaad, 2003; Patil et al., 2012; Swanson et al., 2005; Vasconez et al., 2020).

Weedy Hosts

Amaranthus viridis (slender amaranth)*, *Cyphomandra betaceae* (tamarillo, tree tomato), *Physalis angulata* (cutleaf groundcherry)*, *Portulaca oleracea* (little hogweed)*, *Rumex dentatus* (toothed dock)*, *Solanum dulcamara* (climbing nightshade)*, *Solanum nigrum* (black nightshade)*, *Solanum phureja* (nightshade), and *Urtica dioica* (stinging nettle)* (Ciampi and Sequeira, 1980;Farag et al., 2004; Graham and Lloyd, 1978; Wenneker et al., 1999;Gutarra et al., 2017; Lin et al., 2015; Martin and Nydegger, 1982; Swanepoel, 1992).

Eggplant and pepper can rarely be infected under specific, favorable conditions (Caffier and Hervé, 1996; Martin and French, 1995})

Pest Importance

Ralstonia solanacearum R3 Bvr2 is a serious pathogen with global impacts. According to Elphinstone (2005), about $950 million is lost annually worldwide due to *R. solanacearum* R3 Bvr2.

Ralstonia solanacearum R3 Bvr2 is not known to be present in the United States but could cause significant economic losses by impacting susceptible high-value crops that are grown throughout the country. This pathogen is considered a major problem globally and is among the most destructive pathogens for potatoes (Swanson et al., 2005).

In 2021, potatoes were grown commercially on 933,000 acres in 47 states and the estimated value of the potato harvest was over $3.91 billion and tomatoes were grown on 274,000 acres, with a total value of $1.5 billion. In 2019, geraniums were grown in 47 states with a total value of over $ 217 million (USDA-NASS, 2022).

This pathogen is a select agent in the United States (Janse, 2004; Lambert, 2002; USDA-APHIS-PPQ, 2022). A select agent is a biological agent that has the potential to pose a severe threat to plant health (USDA-APHIS-PPQ, 2022). *R. solanacearum* R3 Bvr2 is also listed as a harmful organism in Canada, El Salvador, Jordan, Mexico, Panama, and Taiwan (APHIS, 2023). In addition, *R. solanacearum* is listed as a quarantine pest in EPPO A2 list (EPPO, 2022a). There would likely be serious trade implications with these countries if this pathogen becomes established in the United States.

Once R3 Bvr2 is introduced and established, it is very difficult to control (Champoiseau et al., 2009) and its lethality, ability to cause latent infections, and environmental survival have made this pathogen a serious problem where it occurs (Gorissen et al., 2004). Eradication efforts are costly, including one eradication effort in a greenhouse in the United States that cost the grower approximately $10 million (Swanson et al., 2007). Control measures are, therefore, aimed at preventing introduction and establishment of the pathogen through exclusion and early detection.

Known Vectors (or associated insects)

There is currently no evidence of *R. solanacearum* being vectored by insects.

Known Distribution

Ralstonia solanacearum R3 Bvr2 has a global distribution and is found in both temperate and tropical climates (Swanson et al., 2005).

Africa: Burundi, Cameroon, Egypt, Ethiopia, Kenya, Madagascar, Malawi, Mauritius, Nigeria, Réunion Island, Rwanda, South Africa, Tanzania, and Uganda. **Asia:** Bangladesh, China, India, Indonesia, Iran, Israel, Japan, Korea, Nepal, Pakistan, Philippines, Sri Lanka, Thailand, Taiwan, and Turkiye. **Central America:** Costa Rica, Guadeloupe, Guatemala. **Europe:** France[†], Georgia, Greece, Hungary, Netherlands[*], Portugal, Russia, Serbia, Spain, and United Kingdom[†]. **North America:** Mexico.

[*] Under eradication

Oceania: Australia and New Guinea. **South America:** Argentina, Bolivia, Brazil, Chile, Colombia, French Guiana, Peru, Uruguay, and Venezuela (Autrique and Potts, 1987; Caruso et al., 2003; Castillo and Plata, 2016; Cellier et al., 2012; Chakraborty and Roy, 2016; Cook and Sequeira, 1991; Cruz et al., 2012; Deberdt et al., 2014; Elphinstone, 2001; Farag et al., 2004; French, 1988; Gabriel et al., 2006; Garcia et al., 1999; Guidot et al., 2009; Hemelda et al., 2019; Horita and Tsuchiya, 2001; Izadiyan and Taghavi, 2013; Janse, 2012; Janse et al., 2004; Jeong et al., 2007; Junaid and Ahmad, 2018; Khoodoo et al., 2010; Lemessa and Zeller, 2007; Mahbou Somo Toukam et al., 2009; Marković et al., 2021; Martin and French, 1995; Matveeva et al., 2003; Mwankemwa, 2015; Natural et al., 2005; Nemeth et al., 2002; Nicole et al., 1998; Parkinson et al., 2013; Perea Soto et al., 2011; Popoola et al., 2015; Pradhanang et al., 2000; Prior and Steva, 1990; Ravelomanantsoa et al., 2017; Sagar et al., 2013; Sedighian et al., 2020; Swanson et al., 2007; Thammakijjawat et al., 2001; Tomlinson and Guntber, 1986; Tusiime et al., 1998; Ustun et al., 2009; Uwamahoro et al., 2018; Van Der Wolf et al., 1998; Vasconez et al., 2020; Wang et al., 2017; Wu et al., 2011; Zayamba Kagona, 2008).

Presence in Nurseries/Greenhouses:
Europe: Belgium[†] and Germany[†] (Marco-Noales et al., 2008), and **North America:** Canada[†] (IPPC, 2020; Janse et al., 2004; Roman-Reyna et al., 2021).

Eradication
R. solanacearum R3 Bvr2 has been eradicated in the United States, Lebanon, and Sweden (Elia et al., 2017; Persson, 1998).

There are also miscellaneous records for Libya and Zambia, but these could not be verified.

Pathway

Ralstonia solanacearum R3 Bvr2 is primarily water and soilborne and can be spread in contaminated irrigation, runoff, and surface water; equipment; and soil (Caruso et al., 2003; Janse et al., 2004; Milling et al., 2009; Wenneker et al., 1999). It can spread naturally from infected roots to healthy roots of neighboring plants (Autrique and Potts, 1987), and insect and nematode damage to healthy roots can facilitate this transmission (Champoiseau et al., 2009). Transmission does not occur by leaf to leaf contact or water splashing (Swanson et al., 2005).

The pathogen is also readily disseminated through infected propagative material, including potato tubers (Wenneker et al., 1999) and geranium cuttings (Milling et al., 2009). Infected plants may remain symptomless but still shed bacteria through their roots, making runoff water an important source of infection (Swanson et al., 2005). Latently infected geranium cuttings were responsible for introduction of *R. solanacearum* R3 Bvr2 in the United States, although it was subsequently eradicated (Dickstein et al., 2017). *R. solanacearum* R3 Bvr2 persists in field soil in the absence of host plants (Elphinstone, 1996; Stander et al., 2003) and remains viable in the roots of one alternative weed host, *Solanum dulcamara*, for many years (Elphinstone, 2001).

Use the PPQ Commodity Import and Export manuals listed below to determine 1) if host plants or material are allowed to enter the United States from countries where the organism is present and 2) what phytosanitary measures (e.g., inspections, phytosanitary certificates, post entry quarantines, mandatory treatments) are in use. These manuals are updated regularly.

Agricultural Commodity Import Requirements (ACIR): ACIR provides a single source to search for and retrieve entry requirements for imported commodities.

Plants for Planting Manual: This manual is a resource for regulating imported plants or plant parts for propagation, including buds, bulbs, corms, cuttings, layers, pollen, scions, seeds, tissue, tubers, and like structures.

Treatment Manual: This manual provides information about treatments applied to imported and domestic commodities to limit the movement of agricultural pests into or within the United States.

Potential Distribution within the United States

Ralstonia solanacearum R3 Bvr2 is not known to be established in the United States due to successful eradication following detections in greenhouses in several states from imported geranium cuttings (Dickstein et al., 2017; USDA-APHIS-PPQ, 2020; Williamson et al., 2002; Roman-Reyna et al., 2021; Strider et al., 1981).

Nearly every state grows at least one of the major crops that can host *R. solanacearum* R3 Bvr2 and all solanaceous crop growing areas may be at risk due to the cold tolerance of *R. solanacearum* R3 Bvr2. However, R3 Bvr2 in infected geranium or potato tuber tissue would not survive the repeated freeze-thaw cycles common in Colorado, Idaho, Maine, Michigan, North Dakota, and Wisconsin (Scherf et al. 2010).

The majority of potatoes are harvested in California, Colorado, Florida, Idaho, Maine, Michigan, Minnesota, North Dakota, Oregon, Washington, and Wisconsin (USDA-NASS, 2022), while California and Florida are the major producers of tomatoes (AgMRC, 2021). In addition, geraniums are popular landscaping plants and widely grown and distributed in the United States. Michigan, California, New Jersey, Pennsylvania, North Carolina, New York, Ohio, Connecticut, Florida, and Wisconsin are major producers of geraniums (USDA-NASS, 2020).

Survey and Key Diagnostics

For the current approved methods and guidance for survey and identification, see Approved Methods for Pest Surveillance (AMPS) pest page on the CAPS Resource and Collaboration website.

References

AgMRC. 2021. Tomatoes. Agricultural Marketing Resource Center. https://www.agmrc.org/commodities-

products/vegetables/tomatoes#:~:text=In%202020%2C%20approximately%2012%2C619.2%20tons,value%20of%20approximately%20%241%20billion.
APHIS. 2023. Phytosanitary Export Database (PExD). United States Department of Agriculture, Animal and Plant Health Inspection Service (APHIS). https://pcit.aphis.usda.gov/pcit/faces/signIn.jsf
Autrique, A., and M. J. Potts. 1987. The influence of mixed cropping on the control of potato bacterial wilt (*Pseudomonas solanacearum*). Annals of Applied Biology 111(1):125-133.
Caffier, D., and A. Hervé. 1996. La pourriture brune de la pomme de terre, un nouveau risque bactérien en Europe [Brown rot of potatoes, a new bacterial risk in Europe]. La Pomme de Terre Française 492:20-23.
Caruso, P., E. Bertonlini, M. Cambra, and M. M. Lopez. 2003. A new and sensitive co-operational polymerase chain reaction for rapid detection of *Ralstonia solanacearum* in water. Journal of Microbiological Methods 55(1):257-272.
Castillo, J. A., and G. Plata. 2016. The expansion of brown rot disease throughout Bolivia: Possible role of climate change. Canadian Journal of Microbiology 62(5):442-448.
Cellier, G., B. Remenant, F. Chiroleu, P. Lefeuvre, and P. Prior. 2012. Phylogeny and population structure of brown rot and Moko disease-causing strains of *Ralstonia solanacearum* phylotype II. Applied and Environmental Microbiology:2367–2375.
Chakraborty, R., and T. S. Roy. 2016. Threats faced by brown rot of potato in Bangladesh. Microbiology Research 7(1):8-12.
Champoiseau, P. G., J. B. Jones, and C. Allen. 2009. *Ralstonia solanacearum* race 3 biovar 2 causes tropical losses and temperate anxieties. Plant Health Progress DOI:10.1094/PHP-2009-0313-01-RV.
Champoiseau, P. G., and M. Timur. 2009. USDA-NRI Project: *R. solanacearum* race 3 biovar 2: detection, exclusion and analysis of a select agent educational modules: bacterial wilt of tomato.
Ciampi, L., and L. Sequeira. 1980. Influence of temperature on virulence of race 3 strains of *Pseudomonas solanacearum*. American Potato Journal 57(7):307-317.
Cook, D., and L. Sequeira. 1991. The use of subtractive hybridization to obtain a DNA probe specific for *Pseudomonas solanacearum* race 3. Molecular and General Genetics 227(3):401-410.
Cook, J. C., and J. Floyd. 2020. New Pest Response Guidelines: *Ralstonia solanacearum* (Smith, 1896) Yabuuchi et al., 1996 “race 3 biovar 2”. USDA–APHIS–PPQ–Emergency and Domestic Programs, Riverdale, Maryland. 52 pp.
Cruz, L., M. Eloy, F. Quirino, H. Oliveira, and R. Tenreiro. 2012. Molecular epidemiology of *Ralstonia solanacearum* strains from plants and environmental sources in Portugal. European Journal of Plant Pathology 133(3):687-706.
Deberdt, P., J. Guyot, R. Coranson-Beaudu, J. Launay, M. Noreskal, P. Rivière, F. Vigné, D. Laplace, L. Lebreton, and E. Wicker. 2014. Diversity of *Ralstonia solanacearum* in French Guiana expands knowledge of the “emerging ecotype”. Phytopathology 104(6):586-596.
Dickstein, E. R., P. G. Champoiseau, J. B. Jones, T. M. Momol, P. Ji, D. J. Norman, Q. Huang, C. Allen, S. A. Miller, T. Schubert, R. Bulluck, and J. E. Fajardo. 2017. Recovery plan for *Ralstonia solanacearum* race 3 biovar 2 causing brown rot of

potato, bacterial wilt of tomato, and southern wilt of geranium. United States Department of Agriculture, Animal and Plant Health Inspection Service, Agriculture Research Service, National Plant Disease Recovery System. 1-48 pp.
Elia, C., J. Fouad, W. Samer, E. K. Michel.issa, V. Franco, D. Nino, G. Daniele, H. Rosine, A. K. L. Khalil, and S. Emilio. 2017. Surveys of potato-growing areas and surface water in Lebanon for potato brown and ring rot pathogens. Phytopathologia Mediterranea 56(1).
Elphinstone, J. G. 1996. Survival and possibilities for extinction of *Pseudomonas solanacearum* (Smith) Smith in cool climates. Potato Research 39(3):403-410.
Elphinstone, J. G. 2001. Monitoring and control of the potato brown rot bacterium (*Ralstonia solanacearum*) in the UK: A case study. *in* Proceedings of the FNK/EAPR/ESA/UEITP 2nd European Potato Processing Conference., 14-15 November 2001, Lausanne, Switzerland.
Elphinstone, J. G. 2005. The current bacterial wilt situation: A global overview. Pages 9-28 *in* C. Allen, P. Prior, and A. C. Hayward, (eds.). Bacterial Wilt Disease and the Ralstonia solanacearum Species Complex. APS Press, St. Paul, MN.
EPPO. 2022a. Pests recently added to the EPPO A1/A2 Lists or of urgent phytosanitary concern. European and Mediterranean Plant Protection Organization.
EPPO. 2022b. PM 7/21 (3) *Ralstonia solanacearum*, *R. pseudosolanacearum* and *R. syzygii* (*Ralstonia solanacearum* species complex). EPPO Bulletin 52(2):225-261.
Farag, N. S., W. E. Eweda, M. I. Mostafa, and N. M. Balabel. 2004. Phytopathological note: Preliminary observations on the bacteriology and pathology of *Ralstonia solanacearum*. Egyptian Journal of Agricultural Research 82(4):1519-1523.
French, E. R. 1988. Strategies for integrated control of bacterial wilt of potatoes. Pages 133-141 *in* A. C. Hayward and G. L. Hartman, (eds.). Bacterial diseases of the potato: Report of the Planning Conference on Bacterial Diseases of the Potato, 1987. International Potato Center.
Gabriel, D. W., C. Allen, M. Schell, T. P. Denny, J. T. Greenberg, and Y. P. Duan. 2006. Identification of open reading frames unique to a select agent: *Ralstonia solanacearum* race 3 biovar 2. Molecular Plant-Microbe Interactions 19(1):69-79.
Garcia, R., A. Garcia, and L. Delgado. 1999. Distribution, incidence and variability of *Ralstonia solanacearum*, causal agent of bacterial wilt of potato in Merida state, Venezuela. Bioagro 11(1):12-23.
Gorissen, A., S. van Overbeek, and J. D. van Elsas. 2004. Pig slurry reduces the survival of *Ralstonia solanacearum* biovar 2 in soil. Canadian Journal of Microbiology 50(8):587-593.
Graham, J., and A. Lloyd. 1979. Survival of potato strain (race 3) of *Pseudomonas solanacearum* in the deeper soil layers. Australian Journal of Agricultural Research 30(3):489-496.
Graham, J., and A. B. Lloyd. 1978. *Solanum cinereum* R. Br., a wild host of *Pseudomonas solanacearum* biotype II. Journal of the Australian Institute of Agricultural Science 44(2):124-126.
Granada, G. A., and L. Sequeira. 1983. Survival of *Pseudomonas solanacearum* in soil, rhizosphere, and plant roots. Canadian Journal of Microbiology 29:433-440.

Guidot, A., M. Elbaz, S. Carrère, M. I. Siri, M. J. Pianzzola, and P. Prior. 2009. Specific genes from the potato brown rot strains of *Ralstonia solanacearum* and their potential use for strain detection. Phytopathology 99(9):1105-1112.

Gutarra, L., J. Herrera, E. Fernandez, J. Kreuze, and H. Lindqvist-Kreuze. 2017. Diversity, pathogenicity, and current occurrence of bacterial wilt bacterium *Ralstonia solanacearum* in Peru. Frontiers in Plant Science 8:1221. doi: 10.3389/fpls.2017.0122.

Hemelda, N. M., R. Safitri, and S. Suhandono. 2019. Genetic diversity of *Ralstonia solanacearum*, A phytopathogenic bacterium infecting horticultural plants in Java, Indonesia. Biodiversitas 20(2):364-372.

Horita, M., and K. Tsuchiya. 2001. Genetic diversity of Japanese strains of *Ralstonia solanacearum*. Phytopathology 91(4):399-407.

Huerta, A. I., A. Milling, and C. Allen. 2015. Tropical strains of *Ralstonia solanacearum* outcompete race 3 biovar 2 strains at lowland tropical temperatures. Applied and Environmental Microbiology 81(10):3542-3551.

IPPC. 2020. *Ralstonia solanacearum* race 3 biovar 2: detection in a Canada greenhouse / *Ralstonia solanacearum* race 3 biovar 2: Détection dans une serre canadienne (CAN-54/2). International Plant Proection Convention.

Izadiyan, M., and S. M. Taghavi. 2013. Host range variation and genetic diversity of Iranian isolates of *Ralstonia solanacearum* from potato and tomato with RAPD and (GTG) 5-PCR. Journal of Plant Pathology 95(1):87-97.

Janse, J. D. 1996. Potato brown rot in western Europe – history, present occurrence and some remarks on possible origin, epidemiology and control strategies. EPPO Bulletin 26(3-4):679-695.

Janse, J. D. 2004. EPPO Standards: Diagnostic protocols for regulated pests: *Ralstonia solanacearum* (PM 7/21(1)). Bulletin OEPP/EPPO Bulletin 34(2):173-178.

Janse, J. D. 2012. Review on brown rot (*Ralstonia solanacearum* Race 3, Biovar 2, Phylotype IIB) epidemiology and control in the Netherlands since 1995: A success story of integrated pest management. Journal of Plant Pathology 94(2):257-272.

Janse, J. D., H. E. van den Beld, J. Elphinstone, S. Simpkins, N. N. A. Tjou-Tam-Sin, and J. van Vaerenbergh. 2004. Introduction to Europe of *Ralstonia solanacearum* biovar 2, race 3 in *Pelargonium zonale* cuttings. Journal of Plant Pathology 86(2):147-155.

Jeong, Y., J. Kim, Y. Kang, S. Lee, and I. Hwang. 2007. Genetic diversity and distribution of Korean isolates of *Ralstonia solanacearum*. Plant Disease 91(10):1277-1287.

Junaid, M., and M. Ahmad. 2018. Race and biovar determination of *Ralstonia solanacearum* in the north west of Pakistan. Novel Research in Microbiology Journal 2(6):138-146.

Khoodoo, M. H. R., E. S. Ganoo, and A. S. Saumtally. 2010. Molecular characterization and epidemiology of *Ralstonia solanacearum* Race 3 biovar 2 causing brown rot of potato in Mauritius. Journal of phytopathology 158(7-8):503-512.

Lambert, C. 2002. Agricultural Bioterrorism Protection Act of 2002: possession, use, and transfer of biological agents and toxins; Interim and Final Rule (7CRF part

331). Pages 38 *in*. Department of Agriculture Animal and Plant Health Inspection Service.
Lemessa, F., and W. Zeller. 2007. Isolation and characterisation of *Ralstonia solanacearum* strains from Solanaceae crops in Ethiopia. Journal of Basic Microbiology 47(1):40-49.
Lin, C.-H., J.-F. Wang, Y.-F. Wu, and A.-H. Cheng. 2015. Phylotype II race 3 biovar 2 strain of *Ralstonia solanacearum* in Taiwan: Survival potential and virulence on tomato, eggplant, and pepper. Acta Horticulturae 1069:293-300.
Mahbou Somo Toukam, G., G. Cellier, E. Wicker, C. Guilbaud, R. Kahane, and C. Allen. 2009. Broad diversity of *Ralstonia solanacearum* strains in Cameroon. Plant Disease 93(11):1123-1130.
Marco-Noales, E., E. Bertolini, C. Morente, and M. M. López. 2008. Integrated approach for detection of nonculturable cells of *Ralstonia solanacearum* in asymptomatic *Pelargonium* spp. cuttings. Phytopathology 98(8):949-955.
Marković, S., S. Stanković, R. Iličić, S. Veljović Jovanović, S. Milić Komić, A. Jelušić, and T. Popović. 2021. *Ralstonia solanacearum* as a potato pathogen in Serbia: Characterization of strains and influence on peroxidase activity in tubers. Plant Pathology 70(8):1945-1959.
Martin, C., and E. R. French. 1995. Covered-field host range test for *Pseudomonas solanacearum* race 3/biovar 2. Bacterial Wilt Newsletter 12:9.
Martin, C., and U. Nydegger. 1982. Susceptibility of *Cyphomandra betacea* to *Pseudomonas solanacearum*. Plant Disease 66:1025-1027.
Matveeva, E. V., E. S. Pekhtereva, E. V. Nikolaeva, and N. W. Schaad. 2003. Pathogenicity, virulence, and phenotypic diversity of *Ralstonia solanacearum* strains in potato in Russian Federation. Phytopathology 96(6):S58.
Milling, A., F. Meng, T. P. Denny, and C. Allen. 2009. Interactions with hosts at cool temperatures, not cold tolerance, explain the unique epidemiology of *Ralstonia solanacearum* race 3 biovar 2. Phytopathology 99(10):1127-1134.
Moorman, G. W. 2014. Bacterial blight of geranium. Pennsylvania State University. Last accessed 2/2/2023, https://extension.psu.edu/bacterial-blight-of-geranium.
Mwankemwa, Z. 2015. Occurrence and distribution of potato bacterial wilt disease and variability of its causal agent in Southern Highlands of Tanzania. Master of Science, Sokoine University of Agriculture, Morogoro, Tanzania. 60 pp.
Natural, M. P., L. A. D. Lando, and E. M. Jover. 2005. Variability of the potato bacterial wilt pathogen, *Ralstonia solanacearum* (E. F. Smith) Yabuuchi et al., in the Philippines. Pages 493-499 *in* C. Allen, P. Prior, and A. C. Hayward, (eds.). Bacterial wilt disease and the *Ralstonia solanacearum* species complex. APS Press, St. Paul, MN.
Nemeth, J., A. Kovacs, G. Galambos, F. Mero, and M. Molnar. 2002. Occurence of bacterial brown rot of potato *Ralstonia* (*Pseudomonas*) *solanacearum* in Hungary. Novenyvedelem Hungary 38(9):453-461.
Nicole, J. F., J.-J. Cheron, J.-C. Girard, and J. Luisetti. 1998. A tentative explanation of the distribution, on Reunion Island, of bacterial wilt caused by either biovar 2 or biovar 3 of *Ralstonia solanacearum*. Pages 343-350 *in* P. Prior, C. Allen, and J. Elphinstone, (eds.). Bacterial Wilt Disease. Springer, Berlin.

Ozakman, M., and N. W. Schaad. 2003. A real-time BIO-PCR assay for detection of *Ralstonia solanacearum* race 3, biovar 2, in asymptomatic potato tubers. Canadian Journal of Plant Pathology 25(3):232-239.

Parkinson, N., R. Bryant, J. Bew, C. Conyers, R. Stones, M. Alcock, and J. Elphinstone. 2013. Application of variable-number tandem-repeat typing to discriminate *Ralstonia solanacearum* strains associated with English watercourses and disease outbreaks. Appl Environ Microbiol 79(19):6016-6022.

Patil, V. U., J. Gopal, and B. P. Singh. 2012. Improvement for bacterial wilt resistance in potato by conventional and biotechnological approaches. Agricultural Research 1(4):299-316.

Perea Soto, J. M., R. S. García Estrada, R. Allende Molar, J. A. Carrillo Fasio, J. León Félix, B. Valdez Torres, and F. S. M. López Soto. 2011. Identification of races and biovars of isolated from tomato plants. Revista Mexicana de Fitopatología 29(2):98-108.

Persson, P. 1998. Successful eradication of *Ralstonia solanacearum* from Sweden. EPPO Bulletin 28(1-2):113-119.

Popoola, A. R., S. A. Ganiyu, O. A. Enikuomehin, J. G. Bodunde, O. B. Adedibu, H. A. Durosomo, and O. A. Karunwi. 2015. Isolation and characterization of *Ralstonia solanacearum* causing bacterial wilt of tomato in Nigeria. Nigerian Journal of Biotechnology 29(1):1–10.

Pradhanang, P. M., J. G. Elphinstone, and R. T. V. Fox. 2000. Identification of crop and weed hosts of *Ralstonia solanacearum* biovar 2 in the hills of Nepal. Plant Pathology 49(4):403-413.

Prior, P., F. Ailloud, B. L. Dalsing, B. Remenant, B. Sanchez, and C. Allen. 2016. Genomic and proteomic evidence supporting the division of the plant pathogen *Ralstonia solanacearum* into three species. BMC Genomics 17:90.

Prior, P., and H. Steva. 1990. Characteristics of strains of *Pseudomonas solanacearum* from the French West Indies. Plant Disease 74:13-17.

Ravelomanantsoa, S., C. Vernière, A. Rieux, L. Costet, F. Chiroleu, S. Arribat, G. Cellier, O. Pruvost, S. Poussier, I. Robène, F. Guérin, and P. Prior. 2017. Molecular epidemiology of bacterial wilt in the Madagascar Highlands caused by Andean (Phylotype IIB-1) and African (Phylotype III) brown rot strains of the *Ralstonia solanacearum* species complex. Front Plant Sci 8:2258.

Roman-Reyna, V., A. Truchon, P. Sharma, F. P. Hand, R. Mazloom, B. A. Vinatzer, J. M. Jacobs, and C. Allen. 2021. Genome resource: *Ralstonia solanacearum* phylotype II sequevar 1 (race 3 biovar 2) strain UW848 from the 2020 U.S. geranium introduction. Plant Disease 105(1):207-208.

Safni, I., I. Cleenwerck, P. De Vos, M. Fegan, L. Sly, and U. Kappler. 2014. Polyphasic taxonomic revision of the *Ralstonia solanacearum* species complex: proposal to emend the descriptions of *Ralstonia solanacearum* and *Ralstonia syzygii* and reclassify current *R. syzygii* strains as *Ralstonia syzygii* subsp. *syzygii* subsp. nov., *R. solanacearum* phylotype IV strains as *Ralstonia syzygii* subsp. *indonesiensis* subsp. nov., banana blood disease bacterium strains as *Ralstonia syzygii* subsp. *celebesensis* subsp. nov. and *R. solanacearum* phylotype I and III strains as *Ralstoniapseudosolanacearum* sp. nov. International Journal of Systematic and Evolutionary Microbiology 64(Pt_9):3087-3103.

Sagar, V., A. K. Somani, R. K. Arora, S. Sharma, S. K. Chakrabarti, S. K. Tiwari, R. Chaturvedi, and B. P. Singh. 2013. Status of bacterial wilt of potato in the Malwa region of Madhya Pradesh in India. Journal of Plant Pathology 95(2):321-328.

Scherf, J. M., A. Milling, and C. Allen. 2010. Moderate temperature fluctuations rapidly reduce the viability of *Ralstonia solanacearum race* 3, biovar 2, in infected geranium, tomato, and potato plants. Applied and Environmental Microbiology 76(21):7061-7067.

Sedighian, N., S. M. Taghavi, H. Hamzehzarghani, J. M. van der Wolf, E. Wicker, and E. Osdaghi. 2020. Potato-infecting *Ralstonia solanacearum* strains in Iran expand knowledge on the global diversity of brown rot ecotype of the pathogen. Phytopathology® 110(10):1647-1656.

Shekhawat, G. S., and M. C. M. Perombelon. 1991. Factors affecting survival in soil and virulence of *Pseudomonas solanacearum* / Faktoren, die das Überleben und die Virulenz von *Pseudomonas solanacearum* beeinflussen. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz / Journal of Plant Diseases and Protection 98(3):258-267.

Stander, E. I. M., P. S. Hammes, and E. A. Beyers. 2003. Survival of *Ralstonia solanacearum* biovar 2 in soil under different cropping systems. South African Journal of Plant and Soil 20(4):176-179.

Stevenson, W. R., R. Loria, G. D. Franc, and D. P. Weingartner (eds.). 2001. Compendium of Potato Diseases, Second edition. APS Press, St. Paul, MN. 106 pp.

Strider, D. L., R. K. Jones, and R. A. Haygood. 1981. Southern bacterial wilt of geranium caused by *Pseudomonas solanacearum*. Plant Disease 65(1):52-53.

Swanepoel, A. E. 1992. Survival of South African strains of biovar 2 and biovar 3 of *Pseudomonas solanacearum* in the roots and stems of weeds. Potato Research 35(3):329-332.

Swanson, J. K., L. Montes, L. Mejia, and C. Allen. 2007. Detection of latent infections of *Ralstonia solanacearum* race 3 biovar 2 in geranium. Plant Disease 91(7):828-834.

Swanson, J. K., J. Yao, J. Tans-Kersten, and C. Allen. 2005. Behavior of *Ralstonia solanacearum* race 3 biovar 2 during latent and active infection of geranium. Phytopathology® 95(2):136-143.

Thammakijjawat, P., N. Thaveechai, W. Kositratana, J. Chunwongse, R. D. Frederick, and N. W. Schaad. 2001. Genetic analysis of *Ralstonia solanacearum* strains from different hosts in Thailand using PCR-Restriction Fragment Length Polymorphism. Kasetsart Journal (Natural Science) 35:397-408.

Tomlinson, D. L., and M. T. Guntber. 1986. Bacterial wilt in Papua New Guinea. Pages 35-39 *in* G. J. Persley and P. A. Batugal, (eds.). Bacterial wilt disease in Asia and the South Pacific: Proceedings of an international workshop held at PCARRD, Los Baños, Philippines, 8-10 October 1985. Australian Centre for International Agricultural Research.

Tran, T. M., J. M. Jacobs, A. I. Huerta, A. Milling, J. Weibel, and C. Allen. 2016. Sensitive, secure detection of race 3 biovar 2 and native US strains of *Ralstonia solanacearum*. Plant Disease 100(3):630-639.

Tusiime, G., E. Adipala, F. Opio, and A. S. Bhagsari. 1998. Weeds as latent hosts of *Ralstonia solanacearum* in Highland Uganda: Implications to development of an integrated control package for bacterial wilt. Pages 413-419 *in* P. Prior, C. Allen, and J. Elphinstone, (eds.). Bacterial Wilt Disease: Molecular and Ecological Aspects. Springer, Berlin.

USDA-APHIS-PPQ. 2020. Case study: Eradicating *Ralstonia Solanacearum* race 3 biovar 2 from U.S. greenhouses U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine. 18-21 pp.

USDA-APHIS-PPQ. 2022. Select Agents and Toxins List. United States Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine.

USDA-NASS. 2020. 2019 Census of Horticultural Specialties (AC-17-SS-3). National Agricultural Statistics Service, United States Department of Agriculture. 604 pp.

USDA-NASS. 2022. Potatoes 2021 Summary. United States Department of Agriculture National Agricultural Statistics Service (NASS), Washington, DC.

Ustun, N., M. Ozakman, and A. Karahan. 2009. Occurrence of *Ralstonia solanacearum* biovar 2 on tomato, weeds and irrigation water in Turkey. 275-278 pp.

Uwamahoro, F., A. Berlin, C. Bucagu, H. Bylund, and J. Yuen. 2018. Potato bacterial wilt in Rwanda: Occurrence, risk factors, farmers' knowledge and attitudes. Food Security 10(5):1221-1235.

Van Der Wolf, J. M., P. J. M. Bonants, J. J. Smith, M. Hagenaar, E. Nijhuis, J. R. C. M. Van Beckhoven, G. S. Saddler, A. Trigalet, and R. Feuillade. 1998. Genetic diversity of *Ralstonia solanacearum* race 3 in Western Europe determined by AFLP, RC-PFGE and Rep-PCR. Pages 44-49 *in* P. Prior, C. Allen, and J. Elphinstone, (eds.). Bacterial Wilt Disease: Molecular and Ecological Aspects. Springer Berlin Heidelberg, Berlin, Heidelberg.

van Elsas, J. D., P. Kastelein, P. van Bekkum, J. M. van der Wolf, P. M. de Vries, and L. S. van Overbeek. 2000. Survival o*f Ralstonia solanacearum* biovar 2, the causative agent of potato brown rot, in field and microcosm soils in temperate climates. Phytopathology 90(12):1358-1366.

Vasconez, I. N., X. Besoain, P. Vega-Celedón, M. Valenzuela, and M. Seeger. 2020. First report of bacterial wilt caused by *Ralstonia solanacearum* phylotype IIB sequevar 1 affecting tomato in different regions of Chile. Plant Disease 104(7):2023.

Wang, L., B. Wang, G. Zhao, X. Cai, S. Jabaji, P. Seguin, and H. Chen. 2017. Genetic and pathogenic diversity of *Ralstonia solanacearum* causing potato brown rot in China. American Journal of Potato Research 94(4):403-416.

Wenneker, M., M. S. W. Verdel, R. M. W. Groeneveld, C. Kempenaar, A. R. van Beuningen, and J. D. Janse. 1999. *Ralstonia* (*Pseudomonas*) *solanacearum* race 3 (biovar 2) in surface water and natural weed hosts: first report on stinging nettle (*Urtica dioica*). European Journal of Plant Pathology 105(3):307-315.

Williamson, L., K. Nakaho, B. Hudelson, and C. Allen. 2002. *Ralstonia solanacearum* race 3, biovar 2 strains isolated from geraniumare pathogenic on potato. Plant Disease 86(9):987-991.

Wu, Y. F., C. H. Lin, J. F. Wang, and A. S. Cheng. 2011. Population density of *Ralstonia solanacearum* potato strain, phylotype II/race 3/biovar 2, and incidence of potato

bacterial wilt in fields in Dounan, Yunlin County. Plant Pathology Bulletin 20(3/4):68-77.
Zayamba Kagona, J. D. 2008. The incidence of bacterial wilt (*Ralstonia solanacearum*) in informal potato planting material used by farmers in Dedza and Ntcheu districts of Malawi. Masters of Science, Norwegian University of Life Sciences, Ås, Norway. 73 pp.

USDA-APHIS-PPQ-ST staff developed this datasheet. Cite this document as:

PPQ. 2024. Cooperative Agricultural Pest Survey (CAPS) Pest Datasheet for *Ralstonia solanacearum* Race 3 Biovar 2 (Burkholderiaceae): Potato Brown Wilt. United States Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine (PPQ), Raleigh, NC.

Versions

Apr 2015: Updated to include the eradication from Austria.

Feb 2023:
- Added new name to **Synonyms** section.
- Added note to **Taxonomic Position** section.
- Revised **Known Distribution** section; verified references and added new countries.
- Revised **Known Hosts** section. Verified references and added new hosts.
- Revised **Symptoms, Biology and Ecology, Pathway, Potential Distribution, and Survey Instruction** sections.
- Added **Easily Mistaken Species** section.
- Added **Pest Importance** section.

Aug 2026: removed *Solanum cinereum* from the **Known Hosts** section.

Reviewer(s)

Caitilyn Allen, Ph.D., University of Wisconsin-Madison
David J. Norman, Plant Pathologist, University of Florida Apopka, FL.
Qi Huang, Research Plant Pathologist, Floral and Nursery Plants Research, Washington, DC.